

Light-front field theory of hot and dense quark matter*

Michael Beyer

Fachbereich Physik, Universität Rostock, 18051 Rostock, Germany
email: michael.beyer@uni-rostock.de

Abstract. Extending the concepts of light-front field theory to quantum statistics provides a novel approach towards nuclear matter under extreme conditions. Such conditions exist, e.g., in neutron stars or in the early stage of our universe. They are experimentally expected to occur in heavy ion collisions, e.g., at RHIC and accelerators to be build at GSI and CERN. Light-front field theory is particularly suited, since it is based on a relativistic Hamiltonian approach. It allows us to treat the perturbative as well as the nonperturbative regime of QCD and also correlations that emerge as a field of few-body physics and is important for hadronization. Last but not least the Hamiltonian approach is useful for nonequilibrium processes by utilizing, e.g., the formalism of nonequilibrium statistical operators.

1 Finite temperature and the light-front

At first sight, it seems difficult to define proper temperature T for a system “on the light-front”, because there is no Lorentz transformation to the rest frame of an observer, who is holding the thermometer. Despite the fact that this is formally clarified by now (see Sect. 2), I would like to motivate, how temperature could survive “on the light-front”. Consider the Fermi distribution for the canonical ensemble of a noninteracting gas in the instant form

$$f(k) = [\exp(k_{\text{on}}^0/T) + 1]^{-1}, \quad (1)$$

where $k_{\text{on}}^0 = +\sqrt{m^2 + \mathbf{k}^2}$ is the on-shell energy of a one-particle state with respect to the medium frame. Let the velocity of the medium be given by the time-like vector $u = (u^0, \mathbf{u})$, where $u^2 = 1$. If the medium is at rest with respect to the observer, $u = (1, \mathbf{0})$. For convenience we introduce the momentum of the medium¹ $P = M_0 u$ with $P^2 = M_0^2$. Following Susskind [1] we now assume

*Presented at Light-Cone 2004, Amsterdam, 16 - 20 August

¹This can be done, e.g., before the thermodynamic limit, so that the mass of the noninteracting system M_0 can be achieved by summing all the single-particle energies.

an observer moving with a large velocity v along the z axis in the negative z direction. The one-particle momentum components seen from this moving frame are denoted p^μ . The momentum of the medium measured with respect to this system is $P^\mu = (E, 0, 0, P)$ with $E = \sqrt{M_0^2 + P^2}$. As long as $P < \infty$ the relation between the medium system and the observer's system are given by Lorentz transformations $L(\omega_P)$. For $P \rightarrow \infty$ we introduce the momentum fraction η via $p^3 = \eta P$ and expand E and p_{on}^0 , hence $L(\omega_P)$, up to $\mathcal{O}(P^{-2})$ and finally arrive at (in the limit $P \rightarrow \infty$)

$$k^0 = \frac{1}{2} \left(\eta M_0 + \frac{\mathbf{p}_\perp^2 + m^2}{\eta M_0} \right), \quad k^3 = \frac{1}{2} \left(\eta M_0 - \frac{\mathbf{p}_\perp^2 + m^2}{\eta M_0} \right), \quad \mathbf{k}_\perp = \mathbf{p}_\perp. \quad (2)$$

This form suggests to introduce light-cone components and to identify the longitudinal fraction η with the (kinematically invariant) light-cone momentum fraction $x = p^+/P^+ = k^+/M_0$. After doing so (2) becomes the well known transformation of momentum variables [1],

$$k^0 = \frac{1}{2} (k^+ - k^-), \quad k^3 = \frac{1}{2} (k^+ + k^-), \quad \mathbf{k}_\perp = \mathbf{p}_\perp. \quad (3)$$

Hence the Fermi distribution “on the light-front” is given by

$$f(k) = \left[\exp \left(\frac{k_{\text{on}}^- + k^+}{2T} \right) + 1 \right]^{-1}, \quad (4)$$

where the on-shell condition in (1) is evaluated for the light-front coordinates, viz. $k_{\text{on}}^- = (m^2 + \mathbf{k}_\perp^2)/k^+$. Despite the fact that we have introduced light-front coordinates and done a limiting process $P \rightarrow \infty$ the components of the four vector k^μ that appear in (4) are given with respect to the *medium frame*. Also the value for T is still given with respect to the *medium frame*. This is how temperature is usually defined. The thermometer must be in contact with the medium “for a long time” (sometimes called zero’s law of thermodynamics). No Lorentz transformation is now possible to the rest system of an observer. It is not even necessary to connect the temperature to a rest frame of an “*instant form observer*”. The medium frame is relevant. This reflects precisely the different forms of dynamics first introduced by Dirac [2]. Of course arbitrary Lorentz transformations of the medium frame to different frames are still possible and hence a covariant formalism can be achieved also in the light-front form.

2 Statistical physics in the light-front quantization

A formal framework of covariant calculations at finite temperatures *in instant form* has been given in Ref. [3] in a different context. The grand canonical partition operator is given by

$$Z_G = \exp \{ -(u \cdot P - \mu N)/T \}, \quad (5)$$

where μ the chemical potential. The statistical operator is $\rho_G = Z_G / \text{Tr} Z_G$. It is clear that (1) can be achieved by choosing $u^\nu = (1, \mathbf{0})$, which are the components

of u as seen from the medium frame (and, in instant form, the observer's rest frame). In light-front quantization we give up the connection between observer's rest frame and medium frame. As we have seen in the introduction, giving the components of the velocity in the medium frame suffice. We shall arrive at (4) choosing

$$u^\nu = (u^+, u^-, \mathbf{u}^\perp) = (1, 1, 0, 0). \quad (6)$$

Following notation of Ref. [4] the operators appearing in (5) are given by

$$P^\mu = \int d\omega_+ T^{+\mu}(x), \quad N = \int d\omega_+ J^+(x), \quad (7)$$

where $T^{\nu\mu}(x)$ denotes the energy momentum tensor, $J^\nu(x)$ is the conserved current, and $d\omega_+ = dx_- d^2x_\perp$. With the above choice of u the partition operator can then be written as

$$Z_G = \exp \left\{ -\frac{1}{T} \left(\frac{1}{2} P^- + \frac{1}{2} P^+ - \mu N \right) \right\}. \quad (8)$$

The resulting Fermi distribution functions of particles f^+ and antiparticles f^- are then given by [5, 6]

$$f^\pm(k) = [\exp \{ (u \cdot k_{\text{on}} \mp \mu) / T \} + 1]^{-1} \quad (9)$$

with $k_{\text{on}} = (k_{\text{on}}^-, k^+, \mathbf{k}_\perp)$ and $u \cdot k_{\text{on}} = (k_{\text{on}}^- + k^+) / 2$.

3 Many-body Green functions

To treat hot and dense quantum systems, such as nuclear or quark matter in question here, we use the techniques of many-body Green functions. Reference to textbook treatments is given in [7]. Many-body Green functions, even in the nonrelativistic case, are complicated objects. We organize the equations in a Dyson expansion that leads to a hierarchy of linked cluster equations. The Dyson equation approach to correlations in many-body systems along with several approximation schemes has been developed by Schuck and collaborators. A review and references are given in [8]. In recent years we have systematically applied this approach for finite temperatures up to multi-nucleon clusters (see e.g. [9] and refs therein). Here I give a brief summary and the generalization to the light-front dynamics [5, 6]. In fact, the light-cone quantization is well suited to use the many-body Green function formalism as it utilizes the Fock space expansion [4]. We define a light-cone time-ordered Green function for a given number of particles as follows (retarded and advanced many-body Green functions can be defined accordingly)

$$\begin{aligned} \mathcal{G}_{\alpha\beta}^{x^+-x'^+} &= -i \langle T_+ A_\alpha(x^+) A_\beta^\dagger(x'^+) \rangle \\ &= -i \left(\theta(x^+ - x'^+) \langle A_\beta^\dagger(x'^+) A_\alpha(x^+) \rangle \mp \theta(x'^+ - x^+) \langle A_\beta^\dagger(x'^+) A_\alpha(x^+) \rangle \right). \end{aligned} \quad (10)$$

The average $\langle \dots \rangle$ is taken over the exact ground state and the upper (lower) sign is for fermions (bosons). The operators $A_\alpha(x^+)$ could be build out of any number

of field operators (fermions and/or bosons). The light-cone time dependence of the operators is given in the Heisenberg picture by $A(x^+) = e^{iP_+x^+} A e^{-iP_+x^+}$. For $A(x^+) = \Psi(x^+)$, the free Fermion field, the standard light-front propagator is recovered.

In finite temperature formalism the above definition can be generalized. For a grand canonical ensemble on the light-front, the generalized Heisenberg picture assumes the form

$$A(\tau) = e^{H\tau} A e^{-H\tau},$$

where $H = u \cdot P - \mu N$. Note however that H is not the bare Hamiltonian P_+ , but includes additional terms related to Lagrange constraints. Following the convention of [7] the respective Green function is defined by

$$\mathcal{G}_{\alpha\beta}^{\tau-\tau'} = -\langle T_\tau A_\alpha(\tau) A_\beta^\dagger(\tau') \rangle. \quad (11)$$

The average is now taken over the (equilibrium) grand canonical statistical operator ρ_G , viz. $\langle \dots \rangle = \text{tr}\{\rho_G \dots\}$. For $\tau = ix^+$ (“imaginary time”) the Heisenberg picture is formally recovered. The “true” [7] Heisenberg picture with respect to this Hamiltonian H is given by

$$A(x^+) = e^{iHx^+} A e^{-iHx^+}.$$

Using this operator in (10) defines the “real time” many-body Green function. Dyson equations can be established for both forms [8, 10]. In the light-front formalism the Dyson equations are given by [5]

$$i \frac{\partial}{\partial x^+} \mathcal{G}_{\alpha\beta}^{x^+-x'^+} = \delta(x^+ - x'^+) \langle [A_\alpha, A_\beta^\dagger]_\pm \rangle + \sum_\gamma \int d\bar{x}^+ \mathcal{M}_{\alpha\gamma}^{x^+-\bar{x}^+} \mathcal{G}_{\gamma\beta}^{\bar{x}^+-x'^+}. \quad (12)$$

The mass matrix that appears in (12) is given by

$$\mathcal{M}_{\alpha\beta}^{x^+-x'^+} = \delta(x^+ - x'^+) \mathcal{M}_{0,\alpha\beta}^{x^+} + \mathcal{M}_{r,\alpha\beta}^{x^+-x'^+} \quad (13)$$

$$(\mathcal{M}_0 \mathcal{N})_{\alpha\beta}^{x^+} = \langle [[A_\alpha, H](x^+), A_\beta^\dagger(x^+)]_\pm \rangle \quad (14)$$

$$(\mathcal{M}_r \mathcal{N})_{\alpha\beta}^{x^+-x'^+} = \sum_\gamma \langle T_{x^+} [A_\alpha, H](x^+), [A_\beta^\dagger, H](x'^+) \rangle_{\text{irreducible}} \quad (15)$$

where $\mathcal{N}_{\alpha\beta}^{x^+} = \langle [A_\alpha, A_\beta^\dagger]_\pm(x^+) \rangle$. The first term in (13) is instantaneous and related to the mean field approximation, the second term is the retardation or memory term. This form suggests to first solve the mean field problem (neglecting memory) and evaluate higher order contributions involving $\mathcal{M}_{r,\alpha\beta}^{x^+-x'^+}$ in the mean field basis. In this approximation (12) can be written in the following form

$$\left(i \frac{\partial}{\partial x^+} - \mathcal{M}_0^{x^+} \right) \mathcal{G}^{x^+-x'^+} = \delta(x^+ - x'^+) \mathcal{N}^{x^+}, \quad (16)$$

where $\mathcal{M}, \mathcal{G}$ etc. are understood as matrices acting in the index space of $\alpha, \beta \dots$. As an example I give results for the ideal gas (of particles or quasiparticles). Following (almost) the notation of [4] the Fock space representation of H is given by

$$H = \sum \int dp^+ d^2 p_\perp \left[(u \cdot p - \mu) b^\dagger b + (u \cdot p + \mu) d^\dagger d \right], \quad (17)$$

where the sum is over all quantum numbers relevant for the particles b or antiparticles d . It is now straight forward to evaluate $\mathcal{N}$ and $\mathcal{M}_0$ using free fermion field operators in (10). The resulting single-particle light-cone time ordered Green function (ideal gas) is

$$G(k) = \frac{\gamma k_{\text{on}} + m}{\frac{1}{2}z - \frac{1}{2}k_{\text{on}}^- + i\varepsilon} \frac{\theta(k^+)}{2k^+} (1 - f^+) + \frac{\gamma k_{\text{on}} + m}{\frac{1}{2}z - \frac{1}{2}k_{\text{on}}^- - i\varepsilon} \frac{\theta(k^+)}{2k^+} f^+ \quad (18)$$

$$+ \frac{\gamma k_{\text{on}} + m}{\frac{1}{2}z - \frac{1}{2}k_{\text{on}}^- + i\varepsilon} \frac{\theta(-k^+)}{2k^+} f^- + \frac{\gamma k_{\text{on}} + m}{\frac{1}{2}z - \frac{1}{2}k_{\text{on}}^- - i\varepsilon} \frac{\theta(-k^+)}{2k^+} (1 - f^-).$$

where we used (6) and $\frac{1}{2}z = \frac{1}{2}k^- + \mu - \frac{1}{2}k^+$ for convenience. The ensemble averages needed are

$$\langle b^\dagger b \rangle = f^+, \quad \langle b b^\dagger \rangle = 1 - f^+, \quad \langle d^\dagger d \rangle = f^-, \quad \langle d d^\dagger \rangle = 1 - f^-. \quad (19)$$

Eq. (18) is the light-front generalization of equation (31.38) of [7], which is connected to the Matsubara-Fourier representation via the generalized Lehmann representation [7]. The Matsubara frequencies analytically continued to $\frac{1}{2}z$ are given by [5]

$$\frac{1}{2}k_n^- = \begin{cases} i(2n+1)\pi T + \mu - \frac{1}{2}k^+ & \text{fermions,} \\ i2n\pi T + \mu - \frac{1}{2}k^+ & \text{bosons.} \end{cases} \quad (20)$$

In Matsubara representation the $i\varepsilon$ term can be dropped in (18), hence it is obvious that the Pauli factors $f^\pm$ vanish (which is only valid for the one-particle Green function) and the “imaginary time” Green function sums up to

$$G(k_n) = \frac{\gamma k_{\text{on}} + m}{k_n^2 - m^2} \quad (21)$$

where $k_n = (k_n^-, k^+, \mathbf{k}_\perp)$, which have been given in similar form in [11]. Our light-front in-medium Green function (18) given first in [5] shows no basic difference (up to conventions and phase space) from the ones given later [11]. However, we do have used a different representation more useful for actual few-body calculations. Similar evaluation of two- or three-body Fock states leads to the respective two-particle or three-particle equations for the two- and three-body system embedded in a hot and dense medium. The proper Pauli blocking factors follow straight forwardly.

4 Conclusion and prospects

Generalizing many-body Green functions [7] to the light-front and utilizing a cluster expansion we are able to treat few-body correlations in a hot and dense (i.e. relativistic) plasma. As a model we have used an effective zero range interaction [12]. This approach provides the framework to tackle such intriguing questions how hadrons form during the early plasma phase of the universe, if there is a color superconducting phase, what is the nature of QCD phase transition. Also, because of the Hamiltonian form of the light-front quantization, it is possible to invoke the whole formalism of nonequilibrium statistical physics and

derive Kadanov-Baym or Boltzmann equations that could be used to describe the relativistic transport during a heavy ion collision. It is also possible (and necessary) to go beyond the zero range model and implement real light-front QCD, which has been elaborated in the works by Brodsky, Pauli and Pinsky [4].

Acknowledgment: I would like to thank the organizers for the pleasant and inspiring atmosphere during this meeting. I am grateful to H.J. Weber and T. Frederico for all the fruitful discussions and collaboration on the topic. Also I would like to thank S. Mattiello and S. Strauß for their contributions to this work. This work is supported by Deutsche Forschungsgemeinschaft.

References

1. L. Susskind, Phys. Rev. **165** (1968) 1535; J. B. Kogut and D. E. Soper, Phys. Rev. **D1** (1970) 2901.
2. P. A. M. Dirac, Rev. Mod. Phys. **21** (1949) 392.
3. W. Israel, Annals Phys. **100**, 310 (1976); W. Israel, Physics **106A**, 204 (1981); H. A. Weldon, Phys. Rev. D **26**, 1394 (1982).
4. S. J. Brodsky, H. C. Pauli and S. S. Pinsky, Phys. Rept. **301**, 299 (1998), and refs. therein.
5. M. Beyer, S. Mattiello, T. Frederico and H. J. Weber, Phys. Lett. B **521**, 33 (2001).
6. M. Beyer, S. Mattiello, T. Frederico and H. J. Weber Few Body Syst. **31** (2002) 159; M. Beyer, in Bled Workshops in Physics, eds. B. Golli, M. Rosina, S. Sirca, (Ljubljana 2003) p.11, arXiv:hep-ph/0310324; M. Beyer, S. Mattiello, T. Frederico and H. J. Weber, arXiv:hep-ph/0310222.
7. Fetter A L and Walecka J D 1971, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York), and Dover reprints 2003.
8. J. Dukelsky, G. Röpke, and P. Schuck Nucl. Phys. **A 628** (1998) 17 and refs. therein.
9. M. Beyer, G. Röpke and A. Sedrakian, Phys. Lett. B **376** (1996) 7; M. Beyer, W. Schadow, C. Kuhrtz and G. Röpke, Phys. Rev. C **60** (1999) 034004; M. Beyer, S. A. Sofianos, C. Kuhrtz, G. Röpke and P. Schuck, Phys. Lett. B **488** (2000) 247; S. A. Sofianos and M. Beyer, arXiv:nucl-th/0408073.
10. Beyer M. (1997) “Few-body methods to describe quarks, nucleons, and correlated matter”, habilitation thesis (in German) Rostock 1997.
11. V. S. Alves, A. Das and S. Perez, Phys. Rev. D **66** (2002) 125008; A. N. Kvinikhidze and B. Blankleider, Phys. Rev. D **69** (2004) 125005; J. Raufeisen and S. J. Brodsky, arXiv:hep-th/0408108.
12. S. Mattiello and M. Beyer, these proceedings;
S. Strauß, S. Mattiello and M. Beyer, these proceedings.